arXiv:math/0302087v1 [math.DG] 8 Feb 2003

Ricci flow on compact Kähler manifolds of positive bisectional curvature

Huai-Dong Cao, Bing-Long Chen, and Xi-Ping Zhu

1 The main result

We announce a new proof of the uniform estimate on the curvature of solutions to the Ricci flow on a compact Kähler manifold M^n with positive bisectional curvature.

Given a compact Kähler manifold M^n of complex dimension n and a Kähler metric $\tilde{g} = \sum \tilde{g}_{i\bar{j}}(x) dz^i dz^{\bar{j}}$ on M^n of positive holomorphic bisectional curvature, consider the normalized Kähler-Ricci flow

$$\frac{\partial}{\partial t} g_{i\bar{j}}(x,t) = -R_{i\bar{j}}(x,t) + g_{i\bar{j}}(x,t) \tag{1}$$

on M^n with initial condition $g_{i\bar{j}}(x,0) = \tilde{g}_{i\bar{j}}$. By a result of Bishop-Goldberg [2], we can assume the Kähler class of the metric $\tilde{g}$ satisfies the condition:

$$[\tilde{\omega}] = [\tilde{\Sigma}] = \pi c_1(M^n) \tag{2}$$

where $\tilde{\omega} = (\sqrt{-1}/2)\tilde{g}_{i\bar{j}} dz^i \wedge dz^{\bar{j}}$ and $\tilde{\Sigma} = (\sqrt{-1}/2)\tilde{R}_{i\bar{j}} dz^i \wedge dz^{\bar{j}}$ are the Kähler form, the Ricci form of the metric $\tilde{g}$ respectively, while $c_1(M)$ denotes the first Chern class. Under the normalized initial condition (2), the first author [3] (see also Proposition 1.1 in [4]) showed that the solution $g(x,t) = \sum g_{i\bar{j}}(x,t) dz^i dz^{\bar{j}}$ to the normalized flow (1) exists for all time. Furthermore by the work of Mok [11] (and Bando [1] for $n = 3$), the solution metric $g(x,t)$ is known to have positive bisectional curvature for any $t > 0$. Our main result is on the uniform estimate of the curvature independent of t:

Main Theorem *Let M^n be a compact Kähler manifold of complex dimension n and $\tilde{g} = \sum \tilde{g}_{i\bar{j}}(x) dz^i dz^{\bar{j}}$ be a Kähler metric on M^n of positive holomorphic bisectional curvature satisfying condition (2). Let $g(x,t)$ be the solution to the normalized Kähler-Ricci flow (1) on M^n with the initial condition $g(x,0) = \tilde{g}$. Then $g(x,t)$ is nonsingular. Namely, the curvature of $g(x,t)$ is uniformly bounded independent of t.*

Whether the solution g above is nonsingular or not has been one of the important open problems in the study of Hamilton's Ricci flow on compact Kähler manifolds. For $n = 1$, the main theorem is proved by Richard Hamilton [9] based on his Harnack inequality and entropy estimate for the Gaussian curvature. (see also the further improvements by B. Chow [8] and Hamilton [10]). In [6,7], Chen-Tian obtained the uniform estimate on the curvature of $g(x,t)$ based on a Moser-Trudinger type inequality in [16] for Kähler-Einstein manifolds and some new functionals. Note the existence of Kähler-Einstein metrics on M^n follows from the solution to the Frankel conjecture by Mori [12] and Siu-Yau [15]. In contrast, our proof of the main theorem does not rely on the exsitence of Kähler-Einstein metrics, but instead on first author's Harnack inequality for the scalar curvature $R(x,t)$ which is a consequence of the Li-Yau-Hamilton estimate for the Kähler-Ricci flow [4], and a very recent local injectivity radius estimate of Perelman for the Ricci flow [13]. Thus, this gives a crucial estimate needed in a new proof of the Frankel conjecture via Hamilton's Ricci flow method. In next section, we'll outline the proof of the Main Theorem. Complete details of the proof and discussions about convergence of $g(x,t)$ as $t \to \infty$ will appear elsewhere.

Remark Richard Hamilton and the first author [5] are able to prove the main theorem for $n = 2$ by studying the geometry of singularity model from the blow up limits and certain dimension reduction argument.

2 Outline of the proof

Let $g(x,t)$ be the solution to the normalized Kähler-Ricci flow (1) on compact Kähler manifold M^n with positive bisectional curvature whose initial metric satisfies condition (2). It is easy to see that under flow (1), the volume $V = V(t)$, the total scalar curvature $\int_{M^n} R(x,t)d\nu_t$, and the average scalar curvature

$$r = r(t) = \frac{1}{V}\int_{M^n} R(x,t)d\nu_t$$

of $(M^n, g(x,t))$ are all constant in t. In fact, $r(t) = n$ for all t. Clearly, it suffices to show that the scalar curvature of g is uniformly bounded from above in t:

$$R(x,t) < C,\ \forall x \in M^n, \forall t \in [0,\infty)$$

for some constant $C > 0$ indepedent of t. To do this, let us first recall

The Harnack inequlaity for the scalar curvature (Cao [4]) *For any $x, y \in M^n$ and $0 < t_1 < t_2 < \infty$, the scalar curvature $R(x,t)$ of solution metric $g(x,t)$ satisfies the inequlaity*

$$R(x,t_1) \le R(y,t_2)\frac{e^{t_2}-1}{e^{t_1}-1}e^{\Delta/4}. \tag{3}$$

Here $\Delta = \Delta(x,y,t_1,t_2) = \inf_\gamma \int_{t_1}^{t_2} |\gamma'(\tau)|^2 d\tau$, where the infimum is taken over all space-time curves from (x,t_1) to (y,t_2), and $|\gamma'(\tau)|$ is the velocity of γ at time $\tau \in [t_1,t_2]$.

Now for any $x \in M^n$ and $t > 1$, set $t_1 = t$ and $t_2 = t + 1$. We can find $y \in M^n$ such that $R(y, t+1) = r(t+1) = n$. It then follows from the Harnack inequality (3)

$$\begin{aligned} R(x,t) &\leq R(y,t+1)\frac{e^{t+1}-1}{e^t-1}e^{\Delta(x,y,t,t+1)/4} \\ &\leq n(e+1)e^{\Delta(x,y,t,t+1)/4}. \end{aligned} \tag{4}$$

On the other hand, from (1) one can show that

$$\Delta(x,y,t,t+1) \leq e^2 d^2(x,y;t) \tag{5}$$

where $d(x,y;t)$ denotes the geodesic distance between x and y with respect to the metric $g_{i\bar{j}}(t)$. In particular, by (4) and (5),

$$R(z,t) \leq n(e+1)\exp(\frac{1}{4}e^2)$$

for all $z \in B_t(y,1)$, the geodesic ball centered at y of radius 1 with respect to the metric $g_{i\bar{j}}(t)$. Then by Perelman's no local collapsing result (Theorem 4.1 or its corollary in [13]), there exists a constant $\beta > 0$ independent of t such that the volume of the geodesic ball $B_t(y,1)$ has a uniform lower bound:

$$Vol(B_t(y,1)) \geq \beta. \tag{6}$$

From (6) one can, by a volume comparison argument of Yau (cf. [14]), deduce a uniform upper bound on the diameter d_t of $(M^n, g(x,t))$: there is a constant $D = D(\beta) > 0$ independent of t, such that for all $t > 1$

$$d_t \leq D. \tag{7}$$

But then, (4), (5), and (7) imply that, for any $t > 1$ and any $x \in M^n$,

$$R(x,t) \leq n(e+1)\exp(e^2 D^2/4),$$

which is our desired uniform estimate.

Finally we should remark that the above argument also works for compact Kähler manifolds with nonnegative bisectional curvature.

References

[1] S. Bando, *Compact 3-folds with nonnegative bisectional curvature*, J. Differ. Geom. **19** (1984), 283-297.

[2] R. L. Bishop and S. I. Goldberg, *On the second cohomologh group of a Kaehler manifold of positive curvature*, Proc. Amer. Math. Soc. **16** (1965), 119-122.

[3] H.-D. Cao, *Deformation of Kähler metrics to Kähler -Einstein metrics on compact Kähler manifolds*, Invent. Math. **81** (1985), 359-372.

[4] H.-D. Cao, *On Hanarck's inequalities for the Kähler-Ricci flow*, Invent. Math. **109** (1992), 247-263.

[5] H.-D. Cao and R. S. Hamilton, unpublished work.

[6] X. Chen and G. Tian, *Ricci flow on Kähler-Einstein surfaces*, Invent. Math. **147** (2002), 487-544.

[7] X. Chen and G. Tian, *Ricci flow on Kähler-Einstein manifolds*, preprint.

[8] B. Chow, *The Ricci flow on the* 2*-sphere*, J. Differ. Geom. **33** (1991), 325-334.

[9] R. S. Hamilton, *The Ricci flow on surfaces*, in Mathematics and General Relativity, Contemporary Mathematics **71** (1988), 237-261.

[10] R. S. Hamilton, *An isoperimetric estimate for the Ricci flow on the 2-sphere*, Modern methods in complex analysis (Princeton, NJ, 1992), 201-222, Ann. of Math. Stud. **137**, Princeton Univ. Press, 1995.

[11] N. Mok, *The uniformization theorem for compact Kähler manifolds of nonnegative holomorphic bisectional curvature*, J. Differ. Geom. **27** (1988), 179-214.

[12] S. Mori, *Projective manifolds with ample tangent bundles*, Ann. Math. **76** (1979), 213-234.

[13] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, preprint.

[14] R. Schoen and S.-T. Yau, *Lecture on Differential Geometry*, Conference Proceedings and Lecture Notes in Geometry and Topology, I. International Press, Cambridge, MA, 1994.

[15] Y.-T. Siu and S.-T. Yau, *Compact Kähler manifolds of positive bisectional curvature*, Inven. Math. **59** (1980), 189-204.

[16] G. Tian, *Kähler-Einstein metrics with positive scalar curvature*, Invent. Math. **130** (1997), 1-39.

Institute for Pure and Applied mathematics at UCLA, and Texas A&M University
E-mail: hcao@ipam.ucla.edu or cao@math.tamu.edu

Department of Mathematics, Zhongshang University, & The Institute of Mathematical Sciences, The Chinese University of Hong Kong

Email: blchen@math.cuhk.edu.hk & stszxp@zsu.edu.cn